

Discover a

complete treatment

BRUKINSA[®] (zanubrutinib) is a prescription medicine used to treat adults with:

- Waldenström's macroglobulinemia (WM).
- Mantle cell lymphoma (MCL) who have received at least one prior treatment for their cancer.
- Marginal zone lymphoma (MZL) when the disease has come back or did not respond to treatment and who have received at least one certain type of treatment.

BRUKINSA was approved for MCL and MZL based on response rate. There are ongoing evaluations to confirm clinical benefit for these uses. It is not known if BRUKINSA is safe and effective in children.

IMPORTANT SAFETY INFORMATION

BRUKINSA may cause serious side effects, including: Bleeding problems (hemorrhage) that can be serious and may lead to death. Your risk of bleeding may increase if you are also taking a blood thinner medicine.

Please see additional Important Safety Information throughout, and accompanying full [Patient Information](#).





Why is a BTK inhibitor important for the treatment of WM, MCL, and MZL?

These B-cell lymphomas are caused by the rapid growth and spread of cancerous B cells.

- Bruton’s tyrosine kinase (BTK) is a protein that signals to cancerous B cells, helping them to grow and spread
- Blocking BTK can help stop this signaling

BRUKINSA is a BTK inhibitor that was designed to **completely block BTK.**

BRUKINSA has been shown to block 100% of BTK in blood cells and 94% to 100% of BTK in lymph nodes when taken at the recommended total daily dose of 320 mg.

The significance of completely blocking BTK on treatment responses has not been established.

Please see additional Important Safety Information throughout, and accompanying full [Patient Information](#).



IMPORTANT SAFETY INFORMATION

BRUKINSA may cause serious side effects, including:
Bleeding problems (hemorrhage) (continued). Tell your healthcare provider if you have any signs or symptoms of bleeding, including: blood in your stools or black stools (looks like tar), pink or brown urine, unexpected bleeding, or bleeding that is severe or you cannot control, vomit blood or vomit that looks like coffee grounds, cough up blood or blood clots, increased bruising, dizziness, weakness, confusion, change in speech, or headache that lasts a long time.



Treatment that is

complete
and consistent

IMPORTANT SAFETY INFORMATION

BRUKINSA may cause serious side effects, including:

Infections that can be serious and may lead to death. Tell your healthcare provider right away if you have fever, chills, or flu-like symptoms.

What are the most common side effects?

The most common side effects of BRUKINSA include:

- Decreased white blood cells
- Upper respiratory tract infection
- Decreased platelet count
- Bleeding
- Rash
- Muscle or joint pain

These are not all the possible side effects of BRUKINSA.

Call your doctor for medical advice about side effects.


You may report side effects to FDA at 1-800-FDA-1088.

Please see additional Important Safety Information throughout, and accompanying full Patient Information.



How do I take BRUKINSA?

The recommended dose of BRUKINSA is 320 mg daily, which is four 80-mg capsules.



BRUKINSA capsules should be taken whole with water—do not open, break, or chew. BRUKINSA can be taken with or without food.

BRUKINSA may cause serious side effects, including: Heart rhythm problems (atrial fibrillation and atrial flutter).

Tell your healthcare provider if you have any of the following signs or symptoms: your heartbeat is fast or irregular, you feel lightheaded or dizzy, pass out (faint), have shortness of breath, or have chest discomfort.

How do I know what dosing schedule is right for me?

Your doctor will select a dose schedule that best fits your needs.

For example, if you take other medications twice a day, your doctor may recommend that you take BRUKINSA twice a day to keep your medications on a similar schedule.

How long will I be on treatment with BRUKINSA?

You will take BRUKINSA for as long as your doctor thinks it is helping you, or for as long as side effects are manageable.

Your **dose and schedule may be changed or interrupted** by your doctor to meet your individual treatment needs, including managing side effects.

Please see additional Important Safety Information throughout, and accompanying full Patient Information.



Dosing designed for

c mplete

around-the-clock
treatment

IMPORTANT SAFETY INFORMATION

BRUKINSA may cause serious side effects, including:

Decrease in blood cell counts. Decreased blood counts (white blood cells, platelets, and red blood cells) are common with BRUKINSA, but can also be severe. Your healthcare provider should do blood tests during treatment with BRUKINSA to check your blood counts.

What if I miss a dose?

If you miss a dose of BRUKINSA, take it as soon as you remember on the same day. Return to your normal schedule the next day.

It's important to take BRUKINSA as directed in order to ensure you always have the amount of medicine your healthcare provider thinks is **optimal for you.**

Tell your healthcare provider about all the medicines you take.

Taking BRUKINSA with certain other medications may affect how BRUKINSA works and can cause side effects.

See the next page to review questions about taking other medicines while on BRUKINSA.

Please see additional Important Safety Information throughout, and accompanying full [Patient Information](#).



Can I take other medicines while on BRUKINSA?

Tell your healthcare provider about all the medications you are currently taking, including:

- Prescription and over-the-counter medicines
- Vitamins and herbal supplements

Examples of supplements and foods to inform your doctor about include:

- Echinacea
- Ginseng
- Goldenseal
- St. John's wort
- Grapefruit juice
- Seville oranges

IMPORTANT SAFETY INFORMATION

BRUKINSA may cause serious side effects, including:

Second primary cancers. New cancers have happened in people during treatment with BRUKINSA, including cancers of the skin or other organs. Your healthcare provider will check you for other cancers during treatment with BRUKINSA. Use sun protection when you are outside in sunlight.

What questions should I ask my doctor?

Here are some questions you may find beneficial to ask your doctor and nurse:

- How often will I need to come in for appointments?
- What is the correct way to take my medication?
- How will I know if I am seeing results from treatment?
- What should I do if I notice any side effects?

Please see additional Important Safety Information throughout, and accompanying full Patient Information.



myBeiGene®

Patient Support Program

To meet your individual needs, myBeiGene pairs you with a dedicated Oncology Nurse Advocate who will personalize support for you and your caregivers during your treatment.



Simplify access to BRUKINSA



Educate about BRUKINSA



Provide personalized solutions

Call **1-833-BeiGene** (1-833-234-4363) to talk with an Oncology Nurse Advocate 8 AM–8 PM ET Monday through Friday or visit **BRUKINSA.com**.

Please see Important Safety Information throughout, and accompanying full Patient Information.

